

# Extrapolation of vector valued rearrangement operators

Stefan Geiss <sup>\*</sup>      Paul F.X. Müller

September 4, 2007

## Abstract

Given an injective map  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  between the dyadic intervals of the unit interval  $[0, 1)$ , we study extrapolation properties of the induced rearrangement operator of the Haar system  $\text{Id}_X \otimes T_{p,\tau} : L_{X,0}^p([0, 1)) \rightarrow L_X^p([0, 1))$ , where  $X$  is a Banach space and  $L_{X,0}^p$  the subspace of mean zero random variables. If  $X$  is a UMD-space, then we prove that the property that  $\text{Id}_X \otimes T_{p,\tau}$  is an isomorphism for some  $1 < p \neq 2 < \infty$  extrapolates across the entire scale of  $L_X^q$ -spaces with  $1 < q < \infty$ . In contrast, if only  $\text{Id}_X \otimes T_{p,\tau}$  is bounded and not its inverse, then we show that there can only exist one-sided extrapolation theorems.

2000 Mathematics Subject Classification: 46B07, 46B70, 47B37

## 1 Introduction

In vector valued  $L^p$ -spaces we study rearrangement operators of the system

$$\{h_I/|I|^{1/p} : I \in \mathcal{D}\},$$

where  $\mathcal{D}$  denotes the collection of all dyadic intervals included in  $[0, 1)$  and  $h_I$  is the  $L_\infty$ -normalized Haar function with support  $I$ . These rearrangement

---

<sup>\*</sup>Supported by the project *Stochastic and Harmonic Analysis, Interactions and Applications* of the Academy of Finland.

operators are defined by an injective map  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  as extension of

$$\text{Id}_X \otimes T_{p,\tau} : \sum_{I \in \mathcal{D}} a_I h_I / |I|^{1/p} \rightarrow \sum_{I \in \mathcal{D}} a_I h_{\tau(I)} / |\tau(I)|^{1/p},$$

where  $(a_I)_{I \in \mathcal{D}} \subseteq X$  is finitely supported and  $X$  is a Banach space. This paper continues [11] and is related in spirit to [8]. In particular, we are motivated by extrapolation properties of vector valued martingale transforms, i.e. maps of type

$$\sum_{I \in \mathcal{D}} a_I h_I \rightarrow \sum_{I \in \mathcal{D}} c_I a_I h_I \tag{1}$$

where  $(a_I)_{I \in \mathcal{D}} \subseteq X$  is finitely supported and  $(c_I)_{I \in \mathcal{D}} \in \ell_\infty(\mathcal{D})$ . Extrapolation theorems for these martingale transforms were widely studied in the literature and go back, for example, to Maurey [9] and Burkholder-Gundy [6] (see [5] for a general overview). In our setting these classical theorems state that if (1) is bounded on  $L_X^p$  for some  $p \in (1, \infty)$ , then it is bounded on  $L_X^q$  for all  $q \in (1, \infty)$ . The significance of those theorems can be already seen in the scalar valued setting: Since a martingale transform is trivially bounded on  $L^2$ , extrapolation yields its boundedness on each of the spaces  $L^q$  with  $q \in (1, \infty)$ . The aim of this paper is to analyze the extrapolation properties of the family  $\text{Id}_X \otimes T_{p,\tau}$ .

In Section 3 we start by two examples. Example 3.1 shows that the continuity of a 'typical' permutation  $\text{Id}_X \otimes T_{p,\tau}$  already implies that  $X$  has to have the UMD-property. The second example provides a permutation such that the continuity of  $\text{Id}_X \otimes T_{p,\tau}$  with  $p \in (1, 2]$  implies the type  $p$  property of the Banach space  $X$ . As a consequence we deduce in Corollary 3.3 that one does not have an upwards extrapolation: For  $X = \ell_p$  and  $p \in (1, 2)$  (so that  $X$  is, in particular, a UMD-space) there is a permutation  $\tau$  such that  $\text{Id}_X \otimes T_{p,\tau}$  is continuous, but  $\text{Id}_X \otimes T_{q,\tau}$  fails to be continuous for  $q \in (p, 2]$ .

The natural question arises whether we still have a one-sided extrapolation meaning that the boundedness of  $\text{Id}_X \otimes T_{p,\tau}$  implies that one of  $\text{Id}_X \otimes T_{q,\tau}$  in the case  $1 < q < p < 2$ .

In Section 4 we answer this to the positive for permutations  $\tau$  satisfying the assumption  $|\tau(I)| = |I|$ . The results are formulated in Theorem 4.2 and Corollary 4.3 and proved by transferring Maurey's classical argument [9] to the permutation case via Proposition 4.4. In Corollary 4.3 we extrapolate the boundedness of  $\text{Id}_X \otimes T_{p,\tau}$  for a UMD-space  $X$  and  $p \in (1, 2)$  *downwards* to 1 to the boundedness of  $\text{Id}_X \otimes T_{q,\tau}$  for  $q \in (1, p)$ .

In Section 5 we do not assume anymore the condition  $|\tau(I)| = |I|$ . In Corollaries 5.6 and 5.7 we obtain a one-sided extrapolation as well. By duality Corollary 5.6 yields a two-sided extrapolation in Theorem 5.8: We show for a UMD-space  $X$  that if  $\text{Id}_X \otimes T_{p,\tau}$  is an isomorphism on some  $L_{X,0}^p$  with  $1 < p \neq 2 < \infty$ , then the rearrangement  $\text{Id}_X \otimes T_{q,\tau}$  is an isomorphism on  $L_{X,0}^q$  for each  $q \in (1, \infty)$ . Thus for a UMD-space valued rearrangement the property of being an isomorphism extrapolates across the entire scale of  $L_{X,0}^q$  spaces,  $q \in (1, \infty)$  – just as for martingale transforms or for scalar valued rearrangements  $T_{q,\tau} : L_0^q \rightarrow L_0^q$ , see [11].

The extrapolation properties of *scalar* valued rearrangement operators are a direct consequence of Pisier’s re-norming of  $H^1$ ,

$$\|g\|_{H^1}^{1-\theta} \sim \sup\{\|\sum |g_I|^{1-\theta} |w_I|^\theta h_I\|_{L^p} : \|w\|_{L^2} = 1\},$$

where  $p \in (1, 2)$ ,  $1/p = 1 - (\theta/2)$ ,  $g = \sum g_I h_I$ , and  $w = \sum w_I h_I$ . This well known fact is recorded for instance in [10] and was exploited further in [8]. As Pisier’s re-norming of  $H^1$  uses the lattice structure of  $L^p$ , our analysis of the *vector* valued case circumvents its use and relies instead on combinatorial and geometric properties of  $\tau$  that hold when  $T_{p,\tau}$  is an isomorphism [11].

## 2 Preliminaries

In the following we equip the unit interval  $[0, 1)$  with the Lebesgue measure  $\lambda$ . The set of dyadic intervals of length  $2^{-k}$  is denoted by  $\mathcal{D}_k$ , the set of all dyadic intervals by  $\mathcal{D}$ , and  $\mathcal{F}_k := \sigma(\mathcal{D}_k)$ . Given  $I \in \mathcal{D}$ , we use  $Q(I) := \{K \subseteq I : K \in \mathcal{D}\}$  and  $h_I$  denotes the  $L_\infty$ -normalized Haar function supported on  $I$ . For a Banach space  $X$  we let  $L_X^p = L_X^p([0, 1))$  be the space of all Radon random variables  $f : [0, 1) \rightarrow X$  such that  $\|f\|_{L_X^p}^p := \int_0^1 \|f(t)\|_X^p dt < \infty$  and  $L_{X,0}^p$  be the sub-space of mean zero random variables, where  $L^p = L_{\mathbb{K}}^p([0, 1))$  and  $L_0^p = L_{\mathbb{K},0}^p([0, 1))$  if nothing is said to the contrary with  $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$ . To avoid artificial special cases we assume that the Banach spaces are at least of dimension one.

**Spaces of type and cotype.** Let  $1 \leq p \leq 2 \leq q < \infty$ . A Banach space  $X$  is of *type*  $p$  (*cotype*  $q$ ) provided that there is a constant  $c > 0$  such that

for all  $n = 1, 2, \dots$  and  $a_1, a_2, \dots, a_n \in X$  one has that

$$\left\| \sum_{k=1}^n r_k a_k \right\|_{L_X^p} \leq c \left( \sum_{k=1}^n \|a_k\|_X^p \right)^{\frac{1}{p}} \left( \left( \sum_{k=1}^n \|a_k\|_X^q \right)^{\frac{1}{q}} \leq c \left\| \sum_{k=1}^n r_k a_k \right\|_{L_X^q} \right),$$

where  $r_1, r_2, \dots$  denote independent Bernoulli random variables. We let  $\text{Type}_p(X) := \inf c$  ( $\text{Cotype}_q(X) := \inf c$ ).

**UMD-spaces.** A Banach space  $X$  is called **UMD-space** provided that for some  $p \in (1, \infty)$  (equivalently, for all  $p \in (1, \infty)$ ) there is a constant  $c_p > 0$  such that

$$\sup_{\theta_k \in [-1, 1]} \left\| \sum_{k=1}^n \theta_k d_k \right\|_{L_X^p} \leq c_p \left\| \sum_{k=1}^n d_k \right\|_{L_X^p}$$

for all  $n = 1, 2, \dots$  and all martingale difference sequences  $(d_k)_{k=1}^n \subseteq L_X^1(\mathcal{F}_n)$  with respect to  $(\mathcal{F}_k)_{k=0}^n$ , i.e.  $d_k$  is  $\mathcal{F}_k$ -measurable and  $\mathbb{E}(d_k | \mathcal{F}_{k-1}) = 0$  for  $k = 1, \dots, n$ . The infimum of all possible  $c_p > 0$  is denoted by  $\text{UMD}_p(X)$ .

Using [4, page 12] it follows that  $\text{UMD}_p(X) = \inf d_p$ , where the infimum is taken over all  $d_p > 0$  such that

$$\sup_{\theta_I \in [-1, 1]} \left\| \sum_{I \in \mathcal{D}} \theta_I a_I h_I \right\|_{L_X^p} \leq d_p \left\| \sum_{I \in \mathcal{D}} a_I h_I \right\|_{L_X^p}$$

for all finitely supported  $(a_I)_{I \in \mathcal{D}} \subseteq X$ . An overview about **UMD-spaces** can be found in [5].

**Hardy spaces.** We recall the definition of Hardy spaces we shall use.

**Definition 2.1.** (i) A function  $a \in L_{X,0}^1(\mathcal{F}_N)$ , where  $N \geq 1$ , is called *atom* provided there exists a stopping time  $\nu : \Omega \rightarrow \{+\infty, 0, \dots, N\}$  such that

- (a)  $a_n := \mathbb{E}(a | \mathcal{F}_n) = 0$  on  $\{n \leq \nu\}$  for  $n = 0, \dots, N$ ,
- (b)  $\|a\|_{L_X^\infty} \mathbb{P}(\nu < \infty) \leq 1$ .

(ii) The space  $H_X^{1,at}(\mathcal{F}_N)$  is given by the norm

$$\|f\|_{H_X^{1,at}} := \inf \sum_{k=1}^{\infty} |\mu_k|, \quad f \in L_{X,0}^1(\mathcal{F}_N),$$

where the infimum is taken over all sequences  $(\mu_k)_{k=1}^\infty \subset [0, \infty)$  and atoms  $(a^k)_{k=1}^\infty$  such that  $f = \sum_{k=1}^\infty \mu_k a^k$  in  $L_X^1(\mathcal{F}_N)$ .

(iii) Given  $p \in [1, \infty)$ , the space  $H_X^p(\mathcal{F}_N)$  is given by the norm

$$\|f\|_{H_X^p} := \left( \mathbb{E} \sup_{n=0, \dots, N} \|\mathbb{E}(f|\mathcal{F}_n)\|_X^p \right)^{\frac{1}{p}}, \quad f \in L_{X,0}^p(\mathcal{F}_N).$$

For an atom  $a$  we have that  $a = 0$  on  $\{\nu = \infty\}$ ,  $\text{supp}(a) \subseteq \{\nu < \infty\}$ , and

$$\mathbb{E}\|a\|_X \leq \|a\|_{L_X^\infty} \mathbb{P}(\nu < \infty) \leq 1.$$

The following inequality is well-known (see [2] and [7], cf. [15]):

$$\|f\|_{H_X^1(\mathcal{F}_N)} \leq \|f\|_{H_X^{1,at}(\mathcal{F}_N)} \leq 18\|f\|_{H_X^1(\mathcal{F}_N)}. \quad (2)$$

**Rearrangement operators.** Let  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  be an injective map. Given a Banach space  $X$  and  $p \in [1, \infty)$ , we define the rearrangement operator  $\text{Id}_X \otimes T_{p,\tau}$  on finite linear combinations of Haar functions as

$$\text{Id}_X \otimes T_{p,\tau} : \sum a_I \frac{h_I}{|I|^{1/p}} \rightarrow \sum a_I \frac{h_{\tau(I)}}{|\tau(I)|^{1/p}}, \quad a_I \in X,$$

and let

$$\|\text{Id}_X \otimes T_{p,\tau}\| := \sup \left\{ \left\| \sum_{I \in \mathcal{D}} a_I \frac{h_{\tau(I)}}{|\tau(I)|^{1/p}} \right\|_{L_X^p} : \left\| \sum_{I \in \mathcal{D}} a_I \frac{h_I}{|I|^{1/p}} \right\|_{L_X^p} \leq 1 \right\}$$

where the supremum is taken over all finitely supported  $(a_I)_{I \in \mathcal{D}} \subseteq X$ . In the case  $\|\text{Id}_X \otimes T_{p,\tau}\| < \infty$  we say that  $\text{Id}_X \otimes T_{p,\tau}$  is bounded because it can be continuously extended to  $L_{X,0}^p([0, 1]) \rightarrow L_X^p([0, 1])$ . The dependence on  $p$  of the operator  $T_{p,\tau}$  disappears when the injection  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  satisfies

$$|\tau(I)| = |I|, \quad I \in \mathcal{D},$$

so that we also use  $T_\tau = T_{p,\tau}$ .

**Semenov's condition.** For a non-empty collection  $\mathcal{C}$  of dyadic intervals we let  $\mathcal{C}^* := \bigcup_{I \in \mathcal{C}} I$ . A rearrangement  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  with

$$|\tau(I)| = |I|$$

satisfies *Semenov's condition* if there is a  $\kappa \in [1, \infty)$  such that

$$\sup_{\mathcal{C} \subseteq \mathcal{D}} \frac{|\tau(\mathcal{C})^*|}{|\mathcal{C}^*|} \leq \kappa < \infty. \quad (3)$$

Given  $p \in (1, 2)$ , Semenov's theorem [13, 14] asserts that under the restriction  $|\tau(I)| = |I|$ , condition (3) is equivalent to the boundedness of  $T_\tau : L_0^p([0, 1)) \rightarrow L^p([0, 1))$ .

**Carleson's constant.** For a non-empty collection  $\mathcal{E} \subseteq \mathcal{D}$  the *Carleson constant* is given by

$$\llbracket \mathcal{E} \rrbracket := \sup_{I \in \mathcal{E}} \frac{1}{|I|} \sum_{J \subseteq I, J \in \mathcal{E}} |J|.$$

The Carleson constant is linked to rearrangement operators by the following theorem [11, Theorems 2 and 3]: For a bijection  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  the assertion that for some (all)  $p \in (1, \infty)$  with  $p \neq 2$  one has

$$\| \text{Id}_{\mathbb{K}} \otimes T_{p,\tau} : L_{X,0}^p \rightarrow L_X^p \| \cdot \| \text{Id}_{\mathbb{K}} \otimes T_{p,\tau^{-1}} : L_{X,0}^p \rightarrow L_X^p \| < \infty$$

is equivalent to the existence of an  $A \geq 1$  such that

$$\frac{1}{A} \llbracket \mathcal{E} \rrbracket \leq \llbracket \tau(\mathcal{E}) \rrbracket \leq A \llbracket \mathcal{E} \rrbracket$$

for all non-empty  $\mathcal{E} \subseteq \mathcal{D}$ .

### 3 Two examples

In this section we consider bijections  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  such that  $|\tau(I)| = |I|$  for all  $I \in \mathcal{D}$  and provide examples which show that  $\text{UMD}_p(X)$  and  $\text{Type}_p(X)$  may both be obstructions to the boundedness of

$$\text{Id}_X \otimes T_\tau : L_{X,0}^p \rightarrow L_X^p.$$

From that it becomes clear that Semenov's boundedness criterion [13] does not have a direct correspondence in the vector valued case.

**Example 3.1.** Let  $\tau_0 : \mathcal{D} \rightarrow \mathcal{D}$  be the injection that leaves invariant the intervals of the even numbered dyadic levels. On the odd numbered dyadic levels we define  $\tau_0$  to exchange the dyadic intervals contained in  $[0, 1/2)$  with those contained in  $[1/2, 1)$  by the shifts

$$\tau_0(I) = I + \frac{1}{2} \text{ if } I \subseteq [0, 1/2) \text{ and } \tau_0(I) = I - \frac{1}{2} \text{ if } I \subseteq [1/2, 1).$$

Then one has the following:

- (i) The rearrangement  $\tau_0 = \tau_0^{-1}$  satisfies Semenov's condition with  $\kappa = 2$  so that  $T_{\tau_0}$  is an isomorphism on  $L_0^p$  for  $p \in (1, \infty)$ .
- (ii) For  $p \in (1, \infty)$  one has

$$\frac{1}{3} \text{UMD}_p(X) \leq \| \text{Id}_X \otimes T_{\tau_0} : L_{X,0}^p \rightarrow L_X^p \| \leq 2 \text{UMD}_p(X) \quad (4)$$

so that the boundedness of  $\text{Id}_X \otimes T_{\tau_0}$  on  $L_{X,0}^p$ ,  $p \in (1, \infty)$ , holds precisely when  $X$  satisfies the UMD-property.

PROOF. Assertion (i) is obvious so that let us turn to (ii) and let  $N \geq 2$  be even and recall that  $\mathcal{D}_k$  is the set of dyadic intervals of length  $2^{-k}$ . For  $k \geq 1$  define

$$\mathcal{D}_k^- := \{I \in \mathcal{D}_k : I \subseteq [0, 1/2)\}.$$

The testing functions by which we link the boundedness of  $\text{Id}_X \otimes T_{\tau_0}$  to the UMD-property of  $X$  are

$$f = \sum_{k=1}^N \sum_{I \in \mathcal{D}_k^-} a_I h_I \quad \text{and} \quad g = \sum_{k=1}^{N/2} \sum_{I \in \mathcal{D}_{2k}^-} a_I h_I,$$

where  $a_I \in X$ . Note that  $g$  is obtained from  $f$  by deleting every second dyadic level from the Haar expansion of  $f$  starting with level 1. Consequently,

$$\begin{aligned} \left\| \sum_{k=1}^N (-1)^k \sum_{I \in \mathcal{D}_k^-} a_I h_I \right\|_{L_X^p} &= \|f - 2g\|_{L_X^p} \\ &\leq \|f\|_{L_X^p} + 2\|g\|_{L_X^p} \\ &\leq \|f\|_{L_X^p} + 2\|(\text{Id}_X \otimes T_{\tau_0})f\|_{L_X^p} \end{aligned}$$

$$\begin{aligned}
&\leq (1 + 2\|\text{Id}_X \otimes T_{\tau_0} : L_{X,0}^p \rightarrow L_X^p\|) \|f\|_{L_X^p} \\
&\leq 3\|\text{Id}_X \otimes T_{\tau_0} : L_{X,0}^p \rightarrow L_X^p\| \left\| \sum_{k=1}^N \sum_{I \in \mathcal{D}_k^-} a_I h_I \right\|_{L_X^p}.
\end{aligned}$$

In our definition of  $\text{UMD}_p(X)$  it is sufficient to consider  $\pm 1$  transforms (this is a well-known extreme point argument). Furthermore, by an appropriate augmentation of the filtration we can even restrict ourselves to alternating sequences of signs  $\pm 1$ . Hence we obtain the left hand side of (4) (in fact, we can think to work on  $[0, 1/2)$  as probability space after re-normalization).

For the right hand side of (4) we fix some  $N \geq 1$  and observe that the action of the above rearrangement is an isometry when restricted to  $\sum_{k \text{ odd}, 0 \leq k \leq N} \sum_{I \in \mathcal{D}_k} a_I h_I$  and an isometry when restricted to  $\sum_{k \text{ even}, 0 \leq k \leq N} \sum_{I \in \mathcal{D}_k} a_I h_I$ . Using the  $\text{UMD}$ -property of  $X$ , we merge this information to obtain the boundedness of the rearrangement operator on the entire space  $L_{X,0}^p$ . ■

**Example 3.2.** There exists a rearrangement  $\tau_0 : \mathcal{D} \rightarrow \mathcal{D}$  with  $|\tau_0(I)| = |I|$  satisfying the Semenov condition (3), such that for all  $p \in (1, 2]$  and all Banach spaces  $X$  one has that

$$\text{Type}_p(X) \leq \|\text{Id}_X \otimes T_{\tau_0} : L_{X,0}^p \rightarrow L_X^p\|.$$

PROOF. (a) Fix  $n \geq 1$  and assume disjoint dyadic intervals  $I_0, \dots, I_n$  of the same length, one after each other starting with  $I_0$ . Let

$$\mathcal{A}_k := \{I \in \mathcal{D} : I \subseteq I_k, |I| = 2^{-k}|I_k|\}$$

for  $k = 1, \dots, n$ . We define a permutation  $\tau_n : \mathcal{D} \rightarrow \mathcal{D}$  such that

- (i)  $\mathcal{A}_k$  is shifted from  $I_k$  to  $I_0$  for each  $k = 1, \dots, n$ ,
- (ii) all subintervals of  $I_0$  of length  $2^{-k}|I_0|$ ,  $k = 1, \dots, n$ , are shifted to  $I_1$ ,
- (iii) all subintervals of  $I_1$  of length  $2^{-k}|I_1|$ ,  $k = 2, \dots, n$ , are shifted to  $I_2$ ,
- ...
- (iv) all subintervals of  $I_{n-1}$  of length  $2^{-n}|I_{n-1}|$  are shifted to  $I_n$ .



On all other intervals  $\tau_n$  acts as an identity. One can check that  $\tau_n$  satisfies Semenov's condition with  $\kappa = 3$ . Moreover, for  $a_1, \dots, a_n \in X$ ,

$$\begin{aligned} \int_0^1 \left\| \sum_{k=1}^n \sum_{I \in \mathcal{A}_k} a_k h_I(t) \right\|_X^p dt &= |I_0| \sum_{k=1}^n \|a_k\|_X^p, \\ \int_0^1 \left\| \sum_{k=1}^n \sum_{I \in \mathcal{A}_k} a_k h_{\tau_n(I)}(t) \right\|_X^p dt &= |I_0| \left\| \sum_{k=1}^n r_k a_k \right\|_{L_X^p}^p, \end{aligned}$$

so that

$$\left\| \sum_{k=1}^n r_k a_k \right\|_{L_X^p} \leq \| \text{Id}_X \otimes T_{\tau_n} : L_{X,0}^p \rightarrow L_X^p \| \left( \sum_{k=1}^n \|a_k\|^p \right)^{\frac{1}{p}}$$

where  $r_1, \dots, r_n$  are independent Bernoulli random variables.

(b) Now we 'glue together' the permutations  $\tau_1, \tau_2, \dots$ : to this end we find pairwise disjoint dyadic intervals  $I_0^1, I_1^1 \subseteq [0, 1/2)$ ,  $I_0^2, I_1^2, I_2^2 \subseteq [1/2, 3/4)$ ,  $I_0^3, I_1^3, I_2^3, I_3^3 \subseteq [3/4, 7/8), \dots$ , where  $I_0^n, \dots, I_n^n$  is a collection as in part (a). Defining the permutation  $\tau_0$  on  $I_0^n, \dots, I_n^n$  as in (a) for all  $n = 1, 2, \dots$  and elsewhere as identity, we arrive at our desired permutation  $\tau_0$ . ■

**Corollary 3.3.** *For the permutation  $\tau_0$  from Example 3.2,  $p \in (1, 2)$ , and  $X := \ell_p$  one has*

$$\| \text{Id}_X \otimes T_{\tau_0} : L_{X,0}^p \rightarrow L_X^p \| < \infty$$

but

$$\| \text{Id}_X \otimes T_{\tau_0} : L_{X,0}^q \rightarrow L_X^q \| = \infty \quad \text{for all } q \in (p, 2].$$

PROOF. The first relation follows from Fubini's theorem and the Semenov condition. On the other side,  $X = \ell_p$  is not of type  $q$  as long as  $q \in (p, 2]$  so that  $T_{\tau_0}$  fails to be bounded in  $L_{X,0}^q$ . ■

## 4 Maurey's extrapolation method and the Semenov condition

By Corollary 3.3 we have seen that an extrapolation from  $p$  to  $q$  fails in general if  $q \in (p, 2]$ . Here one should note that the boundedness of  $\text{Id}_X \otimes T_\tau$  :

$L_{X,0}^p \rightarrow L_X^p$  implies the boundedness of  $T_\tau : L_0^p \rightarrow L^p$ , hence the Semenov condition. The aim of this section is to show that, by Maurey's extrapolation method [9], one has an extrapolation from  $p$  to  $q$  in the case that  $q \in (1, p)$ .

**Definition 4.1.** Let  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  be a permutation with  $|\tau(I)| = |I|$ . An operator  $A$  which maps  $f \in L_{X,0}^1(\mathcal{F}_n)$  into a non-negative random variable  $A(f) : [0, 1) \rightarrow [0, \infty)$  and which is homogeneous (i.e.  $A(\mu f) = |\mu|A(f)$ ,  $\lambda$ -a.s., for all  $\mu \in \mathbb{K}$ ), where  $n \geq 1$ , is  $\tau$ -monotone with constant  $c > 0$  provided that one has,  $\lambda$ -a.s., that

$$A\left(\sum_{k=1}^n \gamma_k d_k\right) \leq c \sup_{k=1, \dots, n} |P_{k-1, \tau}(\gamma_k)| A\left(\sum_{k=1}^n d_k\right) \quad (5)$$

for all

$$d_k(t) = \sum_{I \in \mathcal{D}_{k-1}} a_I h_I(t), \quad a_I \in X,$$

and non-decreasing  $(\gamma_k)_{k=1}^n$  with

$$\gamma_k(t) = \sum_{I \in \mathcal{D}_{k-1}} \gamma_k(I) I_I(t), \quad \gamma_k(I) \geq 0,$$

where  $P_{k-1, \tau}(\gamma_k) := \sum_{I \in \mathcal{D}_{k-1}} \gamma_k(I) I_{\tau(I)}(t)$ .

Note that  $P_{k, \tau}(\gamma)$  is correctly defined for all  $\gamma : [0, 1) \rightarrow \mathbb{R}$  that are constant on the dyadic intervals of length  $2^{-k}$ .

**Theorem 4.2.** *For a permutation  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  with  $|\tau(I)| = |I|$  the following assertions are equivalent:*

- (i) *The permutation  $\tau$  satisfies the Semenov condition (3).*
- (ii) *For all  $1 < q < p < \infty$ , Banach spaces  $X$ ,  $n = 1, 2, \dots$ , and  $\tau$ -monotone operators  $A$ , defined on  $L_{X,0}^1(\mathcal{F}_n)$ , with constant  $c > 0$  one has that*

$$\|A : L_{X,0}^q(\mathcal{F}_n) \rightarrow L^q([0, 1))\| \leq d \|A : L_{X,0}^p(\mathcal{F}_n) \rightarrow L^p([0, 1))\|$$

where  $d = d(p, q, c) > 0$  and

$$\|A\|_r = \|A : L_{X,0}^r(\mathcal{F}_n) \rightarrow L^r([0, 1))\| := \sup \left\{ \|A(f)\|_{L^r} : \|f\|_{L_{X,0}^r} \leq 1 \right\}.$$

Before we give the proof of Theorem 4.2 we apply it to our original extrapolation problem.

**Corollary 4.3.** *Let  $X$  be a UMD-space and let  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  be a permutation such that*

$$|\tau(I)| = |I|.$$

*If, for some  $p \in (1, 2)$ , one has that*

$$\text{Id}_X \otimes T_\tau : L_{X,0}^p \rightarrow L_X^p$$

*is bounded, then*

$$\text{Id}_X \otimes T_\tau : L_{X,0}^q \rightarrow L_X^q$$

*is bounded for all  $q \in (1, p)$ .*

PROOF. Because our assumption implies that  $\text{Id}_{\mathbb{K}} \otimes T_\tau : L_0^p \rightarrow L^p$  is bounded it has to satisfy the Semenov condition. We fix  $n \geq 1$  and apply the previous theorem to the operator  $A$  defined, for  $d_k = \sum_{I \in \mathcal{D}_{k-1}} a_I h_I$  with  $a_I \in X$ , as

$$A \left( \sum_{k=1}^n d_k \right) := \int_{\Omega} \left\| (\text{Id}_X \otimes T_\tau) \left( \sum_{k=1}^n r_k(\omega) d_k \right) \right\|_X d\mathbb{P}(\omega)$$

where  $r_1, \dots, r_n$  are independent Bernoulli random variables. It is easy to see that  $A$  satisfies (5) with  $c = 1$ . Moreover by the UMD-property we have

$$\left\| A \left( \sum_{k=1}^n d_k \right) \right\|_{L^p} \sim \left\| (\text{Id}_X \otimes T_\tau) \left( \sum_{k=1}^n d_k \right) \right\|_{L_X^p},$$

where the multiplicative constants do not depend on  $n$ . Hence Theorem 4.2 yields the assertion. ■

The maximal inequality of the following Proposition 4.4 provides the link between rearrangements satisfying Semenov's condition and Maurey's extrapolation technique in [9].

**Proposition 4.4.** *Assume that Semenov's condition (3) is satisfied for a permutation  $\tau$  with  $|\tau(I)| = |I|$  and that  $0 \leq Z_0 \leq Z_1 \leq \dots \leq Z_n$  is a sequence of functions  $Z_k : [0, 1) \rightarrow [0, \infty)$ , where  $Z_k$  is constant on all dyadic intervals of length  $1/2^k$ . Then one has that*

$$\int_0^1 \sup_{k=0, \dots, n} (P_{k, \tau}(Z_k))(t) dt \leq \kappa \int_0^1 Z_n(t) dt.$$

PROOF. Let  $\Delta_0 := Z_0$  and  $\Delta_k := Z_k - Z_{k-1}$  for  $k = 1, \dots, n$ , and let us write

$$\Delta_k = \sum_{I \in \mathcal{D}_k} a_I 1_I$$

with  $a_I \geq 0$ . Fix  $k \in \{0, \dots, n\}$  and observe that, point wise,

$$P_{k, \tau} 1_I \leq 1_{\tau(Q(I))^*} \quad \text{with} \quad Q(I) = \{K \subseteq I : K \in \mathcal{D}\}$$

for  $I \in \mathcal{D}_{k'}$  with  $k' = 0, \dots, k$  (note that  $1_I$  is constant on the dyadic intervals of length  $2^{-k}$  so that we may apply  $P_{k, \tau}$ ). This implies that

$$P_{k, \tau} \left( \sum_{k'=0}^k \sum_{I \in \mathcal{D}_{k'}} a_I 1_I \right) \leq \sum_{k'=0}^k \sum_{I \in \mathcal{D}_{k'}} a_I 1_{\tau(Q(I))^*}.$$

Because the expression on the right-hand side is monotone in  $k$  we conclude that

$$\sup_{k=0, \dots, n} P_{k, \tau} \left( \sum_{k'=0}^k \sum_{I \in \mathcal{D}_{k'}} a_I 1_I \right) \leq \sum_{k'=0}^n \sum_{I \in \mathcal{D}_{k'}} a_I 1_{\tau(Q(I))^*}.$$

Integration gives

$$\int_0^1 \left[ \sup_{k=0, \dots, n} P_{k, \tau} \left( \sum_{k'=0}^k \sum_{I \in \mathcal{D}_{k'}} a_I 1_I \right) (t) \right] dt \leq \sum_{k'=0}^n \sum_{I \in \mathcal{D}_{k'}} a_I |\tau(Q(I))^*|.$$

Our hypothesis gives  $|\tau(Q(I))^*| \leq \kappa |I|$  so that

$$\sum_{k'=0}^n \sum_{I \in \mathcal{D}_{k'}} a_I |\tau(Q(I))^*| \leq \kappa \sum_{k'=0}^n \sum_{I \in \mathcal{D}_{k'}} a_I |I| = \kappa \int_0^1 \left[ \sum_{k=0}^n \Delta_k(t) \right] dt$$

and we are done because

$$\int_0^1 \left[ \sup_{k=0, \dots, n} P_{k, \tau} \left( \sum_{k'=0}^k \sum_{I \in \mathcal{D}_{k'}} a_I 1_I \right) (t) \right] dt = \int_0^1 \sup_{k=0, \dots, n} (P_{k, \tau} Z_k)(t) dt$$

and

$$\int_0^1 \left[ \sum_{k=0}^n \Delta_k(t) \right] dt = \int_0^1 Z_n(t) dt.$$

■

PROOF OF Theorem 4.2. (i)  $\implies$  (ii) We let  $\frac{1}{q} = \frac{1}{r} + \frac{1}{p}$  and

$$d_k := \sum_{I \in \mathcal{D}_{k-1}} \alpha_I h_I \quad \text{so that} \quad T_\tau d_k = \sum_{I \in \mathcal{D}_{k-1}} \alpha_I h_{\tau(I)}.$$

Define  $X_0 := 0$ ,  $X_k := d_1 + \dots + d_k$  for  $k = 1, \dots, n$ ,  $X_k^* := \sup_{l=0, \dots, k} \|X_l\|_X$  for  $k = 0, \dots, n$ ,  ${}^*X_k := X_{k-1}^* + \sup_{l=1, \dots, k} \|d_l\|_X$  for  $k = 1, \dots, n$ ,

$$\gamma_k := ({}^*X_k + \delta)^\alpha$$

for some  $\delta > 0$ ,

$$\alpha := 1 - \frac{q}{p},$$

and

$$\beta_k := P_{k-1, \tau} \gamma_k.$$

By definition we have that

$$\frac{T_\tau(d_k)}{\beta_k} = T_\tau \left( \frac{d_k}{\gamma_k} \right).$$

From the monotonicity assumption on the operator  $A$  it follows that

$$\left\| A \left( \sum_{k=1}^n d_k \right) \right\|_{L^q} \leq c \|\beta_n^*\|_{L^r} \left\| A \left( \sum_{k=1}^n \frac{d_k}{\gamma_k} \right) \right\|_{L^p} \leq c \|A\|_p \|\beta_n^*\|_{L^r} \left\| \sum_{k=1}^n \frac{d_k}{\gamma_k} \right\|_{L_X^p}.$$

From [9, Lemma A] we know that

$$\left\| \sum_{k=1}^n \frac{d_k}{\gamma_k} \right\|_{L_X^p} \leq \frac{p}{q} (\mathbb{E}({}^*X_n + \delta)^q)^{\frac{1}{p}}$$

$$\leq \frac{p}{q} 3^{\frac{q}{p}} (\mathbb{E}(X_n^* + \delta)^q)^{\frac{1}{p}}.$$

Finally, applying Proposition 4.4 we get

$$\begin{aligned} \|\beta_n^*\|_{L^r}^r &= \int_0^1 \sup_{k=1, \dots, n} |(P_{k-1, \tau}(\gamma_k))(t)|^r dt = \int_0^1 \sup_{k=1, \dots, n} (P_{k-1, \tau}(|\gamma_k|^r))(t) dt \\ &\leq \kappa \int_0^1 |\gamma_n(t)|^r dt = \kappa \int_0^1 |^*X_n(t) + \delta|^{\alpha r} dt \leq 3^{\alpha r} \kappa \int_0^1 |X_n^*(t) + \delta|^{\alpha r} dt. \end{aligned}$$

Combining all estimates, we get

$$\left\| A \left( \sum_{k=1}^n d_k \right) \right\|_{L^q} \leq c \|A\|_p 3^\alpha \kappa^{\frac{1}{r}} (\mathbb{E}|X_n^* + \delta|^{\alpha r})^{\frac{1}{r}} \frac{p}{q} 3^{\frac{q}{p}} (\mathbb{E}|X_n^* + \delta|^q)^{\frac{1}{p}}.$$

By  $\delta \downarrow 0$  and Doob's maximal inequality this implies

$$\left\| A \left( \sum_{k=1}^n d_k \right) \right\|_{L^q} \leq c \|A\|_p \frac{3p}{q-1} \kappa^{\frac{1}{r}} \|d_1 + \dots + d_n\|_{L_X^q}.$$

(ii)  $\implies$  (i) We fix  $X = \mathbb{K}$ ,  $n \in \{1, 2, \dots\}$ , and a permutation  $\tau$  with  $|\tau(I)| = |I|$ . Let  $A(\sum_{k=1}^n d_k) := (\sum_{k=1}^n (T_\tau d_k)^2)^{\frac{1}{2}}$  which is  $\tau$ -monotone with constant  $c = 1$ . Clearly,  $\|Af\|_{L^2} = \|f\|_{L^2}$ . If we have an extrapolation to some  $q \in (1, 2)$ , then by the square function inequality the usual permutation operator is bounded in  $L^q$  with a constant not depending on  $n$ , so that by Semenov's theorem [13] condition (3) has to be satisfied. ■

## 5 Extrapolation and the Carleson condition

In this section we consider rearrangement operators induced by bijections  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  that preserves the Carleson packing condition, that is there is an  $A \geq 1$  such that

$$\frac{1}{A} \llbracket \mathcal{E} \rrbracket \leq \llbracket \tau(\mathcal{E}) \rrbracket \leq A \llbracket \mathcal{E} \rrbracket$$

for all non-empty  $\mathcal{E} \subseteq \mathcal{D}$ . In particular, we do not rely anymore on the a-priori hypothesis that  $|\tau(I)| = |I|$ . The corresponding extrapolation results are formulated in Corollary 5.6, Corollary 5.7, and Theorem 5.8, where we

obtain in Corollary 5.7 an alternative proof of Corollary 4.3 that works without  $X$  being a UMD-space. To shorten the notation we let  $\mathcal{D}_0^N := \bigcup_{k=0}^N \mathcal{D}_k$  for  $N \geq 0$ . Because we use complex interpolation we shall assume that all Banach spaces are complex.

We start with a technical condition which ensures a one-sided extrapolation. The condition will be justified by Examples 5.2 and 5.3 below.

**Definition 5.1.** Let  $X$  be a Banach space,  $\tau : \mathcal{D}_0^N \rightarrow \mathcal{D}_0^L$  be an injection,  $\gamma_I > 0$  for  $I \in \mathcal{D}_0^N$ ,  $p \in (1, \infty)$ , and  $\kappa > 0$ . We say that condition  $C(X, p, \kappa)$  is satisfied, provided that for all  $J_0 \in \mathcal{D}_0^N$  there is a decomposition

$$\{I \in \mathcal{D}_0^N : I \subseteq J_0\} = \bigcup_i \mathcal{K}_i,$$

$\mathcal{K}_i \neq \emptyset$ , such that the following is satisfied:

$$(C1) \quad \sum_i |\mathcal{K}_i^*| \leq \kappa |J_0|.$$

$$(C2) \quad \text{For } 1 = \frac{1}{p} + \frac{1}{q} \text{ and}$$

$$\beta_i := \sup \left\{ \left\| \sum_{I \in \mathcal{K}_i} \gamma_I^{\frac{1}{q}} a_I h_I \right\|_{L_X^p}^q : \left\| \sum_{I \in \mathcal{K}_i} a_I h_I \right\|_{L_X^p} = 1 \right\}$$

$$\text{one has that } \sum_i \beta_i |\tau(\mathcal{K}_i)^*| \leq \kappa |J_0|.$$

$$(C3) \quad \text{There exists } p_* \in [p, \infty) \text{ such that}$$

$$\left( \sum_i \left\| \sum_{I \in \mathcal{K}_i} a_I h_I \right\|_{L_X^{p_*}}^{p_*} \right)^{\frac{1}{p_*}} \leq \kappa \left\| \sum_{J_0 \supseteq I \in \mathcal{D}_0^N} a_I h_I \right\|_{L_X^{p_*}}.$$

**Example 5.2.** We assume that  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  with  $|\tau(I)| = |I|$  satisfies the Semenov condition (3) with constant  $\kappa \in [1, \infty)$ , restrict  $\tau$  to  $\tau_N : \mathcal{D}_0^N \rightarrow \mathcal{D}_0^N$ , and take  $\gamma_I = 1$  for all  $I \in \mathcal{D}_0^N$ . Let  $X$  be arbitrary,  $p \in (1, \infty)$ , and  $J_0 \in \mathcal{D}_0^N$ . Because of

$$\left| \bigcup_{J_0 \supseteq I \in \mathcal{D}_0^N} \tau_N(I) \right| \leq \kappa |J_0|$$

we can take

$$\mathcal{K}_1 := \{I \in \mathcal{D}_0^N : I \subseteq J_0\}$$

and conditions (C1), (C2), and (C3) (for any  $p^*$ ) are satisfied with constant  $\kappa$  uniformly in  $N$ .

**Example 5.3.** Let  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  be a bijection and assume that there is an  $A \geq 1$  such that

$$\frac{1}{A} \llbracket \mathcal{E} \rrbracket \leq \llbracket \tau(\mathcal{E}) \rrbracket \leq A \llbracket \mathcal{E} \rrbracket$$

for all non-empty  $\mathcal{E} \subseteq \mathcal{D}$ . Let  $X$  be a UMD-space and  $\gamma_I := |I|/|\tau(I)|$ . As shown in [11, Theorem 1], the permutation  $\sigma = \tau^{-1}$  satisfies the following property P: There exists an  $M > 0$  such that for all dyadic intervals  $J_0 \in \mathcal{D}$  there exists a decomposition as disjoint union

$$\{I \in \mathcal{D} : I \subseteq J_0\} = \sigma(\mathcal{D}) \cap J_0 = \bigcup_i \sigma(\mathcal{L}_i) \cup \bigcup_i \mathcal{E}_i$$

such that

- (1)  $\llbracket \bigcup_i \mathcal{E}_i \rrbracket \leq M$ ,
- (2)  $\sup_{K \in \mathcal{L}_i} \frac{|\sigma(K)|}{|K|} \leq M \frac{|\sigma(\mathcal{L}_i)^*| + |\mathcal{E}_i^*|}{|\mathcal{L}_i^*|}$  for  $\mathcal{L}_i \neq \emptyset$ ,
- (3)  $\sum_i |\sigma(\mathcal{L}_i)^*| \leq M |J_0|$ .

Now we check the counterparts of (C1), (C2), and (C3) for the 'infinite' permutation  $\tau$ .

Condition (C3): As  $X$  is a UMD-space (and therefore super-reflexive) there is a  $p_0 \in [2, \infty)$  such that for all  $p_* \in [p_0, \infty)$  the space  $X$  has cotype  $p_*$ . This cotype and the UMD-property imply (C3) (the constant may depend on  $p_*$ ).

Condition (C1): We write

$$\bigcup_i \mathcal{E}_i = \{\tilde{I}_1, \tilde{I}_2, \dots\} \quad \text{and} \quad \tilde{\mathcal{L}}_j := \{\tau(\tilde{I}_j)\}$$

so that

$$\{I \in \mathcal{D} : I \subseteq J_0\} = \bigcup_i \sigma(\mathcal{L}_i) \cup \bigcup_j \sigma(\tilde{\mathcal{L}}_j) =: \bigcup_i \mathcal{K}_i \cup \bigcup_j \tilde{\mathcal{K}}_j.$$



Now

$$\sum_i |\mathcal{K}_i^*| + \sum_j |\tilde{\mathcal{K}}_j^*| = \sum_i |\sigma(\mathcal{L}_i)^*| + \sum_j |\tilde{I}_j| \leq M|J_0| + \|\bigcup_i \mathcal{E}_i\| |J_0| \leq 2M|J_0|.$$

Condition (C2): let  $p \in (1, \infty)$  be arbitrary and recall that

$$\beta_i = \sup \left\{ \left\| \sum_{I \in \mathcal{K}_i} \gamma_I^{\frac{1}{q}} a_I h_I \right\|_{L_p^X}^q : \left\| \sum_{I \in \mathcal{K}_i} a_I h_I \right\|_{L_p^X} = 1 \right\},$$

where we assume that the sums over  $I$  are finitely supported, and let

$$\tilde{\beta}_j := \sup \left\{ \left\| \sum_{I \in \tilde{\mathcal{K}}_j} \gamma_I^{\frac{1}{q}} a_I h_I \right\|_{L_p^X}^q : \left\| \sum_{I \in \tilde{\mathcal{K}}_j} a_I h_I \right\|_{L_p^X} = 1 \right\} = \gamma_{\tilde{I}_j}.$$

Because  $\gamma_I = |I|/|\tau(I)|$ , the UMD-property of  $X$  gives

$$\beta_i \leq \text{UMD}_p(X)^q \sup_{I \in \mathcal{K}_i} \frac{|I|}{|\tau(I)|}.$$

Since

$$\sup_{I \in \mathcal{K}_i} \frac{|I|}{|\tau(I)|} \leq M \frac{|\mathcal{K}_i^*| + |\mathcal{E}_i^*|}{|\tau(\mathcal{K}_i)^*|}$$

for  $\mathcal{L}_i \neq \emptyset$  we get

$$\begin{aligned} \sum_i \beta_i |\tau(\mathcal{K}_i)^*| &\leq \text{UMD}_p(X)^q \sum_i \sup_{I \in \mathcal{K}_i} \frac{|I|}{|\tau(I)|} |\tau(\mathcal{K}_i)^*| \\ &\leq \text{UMD}_p(X)^q \sum_i M \frac{|\mathcal{K}_i^*| + |\mathcal{E}_i^*|}{|\tau(\mathcal{K}_i)^*|} |\tau(\mathcal{K}_i)^*| \\ &= M \text{UMD}_p(X)^q \sum_i [|\mathcal{K}_i^*| + |\mathcal{E}_i^*|] \\ &\leq 2M^2 \text{UMD}_p(X)^q |J_0|. \end{aligned}$$

In the same way,

$$\sum_j \tilde{\beta}_j |\tau(\tilde{\mathcal{K}}_j)^*| = \sum_j |\tilde{I}_j| \leq M|J_0|.$$

Finally, if we restrict  $\tau$  to  $\tau_N : \mathcal{D}_0^N \rightarrow \mathcal{D}_0^{L_N}$  with  $L_N$  chosen such that  $\tau(\mathcal{D}_0^N) \subseteq \mathcal{D}_0^{L_N}$ , then (C1), (C2), and (C3) are satisfied with the same constant uniformly in  $N$ .

In the following we use the notation

$$L_X^r(\mathcal{D}_0^N) := L_{X,0}^r(\mathcal{F}_{N+1}), \quad H_X^{1,at}(\mathcal{D}_0^N) := H_X^{1,at}(\mathcal{F}_{N+1}),$$

and  $H_X^1(\mathcal{D}_0^N) := H_X^1(\mathcal{F}_{N+1})$  for  $N = 0, 1, \dots$  to avoid a permanent shift in  $N$  because we are working with the sets  $\mathcal{D}_0^N$  rather than with the  $\sigma$ -algebras  $\mathcal{F}_N$ . Now fix Banach spaces  $X$  and  $Y$  and a bounded linear operator  $S : X \rightarrow Y$ , and define the family of operators  $A_p : L_X^p(\mathcal{D}_0^N) \rightarrow L_Y^p(\mathcal{D}_0^L)$  by

$$A_p \left( \sum_{I \in \mathcal{D}_0^N} a_I h_I \right) := \sum_{I \in \mathcal{D}_0^N} S a_I \gamma_I^{\frac{1}{p}} h_{\tau(I)},$$

where  $\gamma_I > 0$ . We aim at extrapolation theorems for this family of operators and extrapolate - under the condition  $C(X, p, \kappa)$  - from  $L^p$  downwards to  $H^1$  in a first step:

**Theorem 5.4.** *If  $p \in (1, \infty)$  and if assumption  $C(X, p, \kappa)$  holds, then*

$$\|A_1 : H_X^1(\mathcal{D}_0^N) \rightarrow H_Y^1(\mathcal{D}_0^L)\| \leq \frac{18p}{p-1} \kappa^{1+\frac{1}{q_*}} \|A_p : L_X^p(\mathcal{D}_0^N) \rightarrow L_Y^p(\mathcal{D}_0^L)\|$$

where  $1 = (1/p_*) + (1/q_*)$  and  $p_*$  is taken from the definition of  $C(X, p, \kappa)$ .

PROOF. Let  $1 = \frac{1}{p} + \frac{1}{q}$  and let  $a \in H_X^{1,at}(\mathcal{D}_0^N)$  be an atom with associated stopping time  $\nu$  (like in Definition 2.1) and assume first that  $\{\nu < \infty\} = J_0 \in \mathcal{D}_0^N$ . For  $J_0$  we choose the sets  $\mathcal{K}_i$  like in Definition 5.1. Moreover, we use

$$D_q a := \sum_{I \in \mathcal{D}_0^N} \gamma_I^{\frac{1}{q}} a_I h_I \quad \text{and} \quad a_i := \sum_{I \in \mathcal{K}_i} a_I h_I$$

for  $a = \sum_{I \in \mathcal{D}_0^N} a_I h_I$  and

$$\beta_i := \sup \left\{ \left\| \sum_{I \in \mathcal{K}_i} \gamma_I^{\frac{1}{q}} a_I h_I \right\|_{L_X^p}^q : \left\| \sum_{I \in \mathcal{K}_i} a_I h_I \right\|_{L_X^p} = 1 \right\}.$$

We get that

$$\begin{aligned}
\|A_1 a\|_{H_Y^1} &\leq \sum_i \|A_1 a_i\|_{H_Y^1} \\
&= \sum_i \|A_p D_q a_i\|_{H_Y^1} \\
&\leq \sum_i |\tau(\mathcal{K}_i)^*|^{\frac{1}{q}} \|A_p D_q a_i\|_{H_Y^p} \\
&\leq \frac{p}{p-1} \sum_i |\tau(\mathcal{K}_i)^*|^{\frac{1}{q}} \|A_p D_q a_i\|_{L_Y^p} \\
&\leq \frac{p}{p-1} \|A_p\| \sum_i |\tau(\mathcal{K}_i)^*|^{\frac{1}{q}} \|D_q a_i\|_{L_X^p} \\
&\leq \frac{p}{p-1} \|A_p\| \sum_i [|\tau(\mathcal{K}_i)^*| \beta_i]^{\frac{1}{q}} \|a_i\|_{L_X^p} \\
&\leq \frac{p}{p-1} \|A_p\| \sum_i [|\tau(\mathcal{K}_i)^*| \beta_i]^{\frac{1}{q}} |\mathcal{K}_i^*|^{\frac{1}{p} - \frac{1}{p_*}} \|a_i\|_{L_X^{p_*}} \\
&\leq \frac{p}{p-1} \|A_p\| \left( \sum_i \left| [|\tau(\mathcal{K}_i)^*| \beta_i]^{\frac{1}{q}} |\mathcal{K}_i^*|^{\frac{1}{p} - \frac{1}{p_*}} \right|^{q_*} \right)^{\frac{1}{q_*}} \\
&\quad \left( \sum_i \|a_i\|_{L_X^{p_*}}^{p_*} \right)^{\frac{1}{p_*}}
\end{aligned}$$

with  $1 = \frac{1}{q_*} + \frac{1}{p_*}$ . Letting  $r := \frac{q}{q_*}$  and  $1 = \frac{1}{r} + \frac{1}{s}$  we obtain that

$$\sum_i \left| [|\tau(\mathcal{K}_i)^*| \beta_i]^{\frac{1}{q}} |\mathcal{K}_i^*|^{\frac{1}{p} - \frac{1}{p_*}} \right|^{q_*} \leq \left( \sum_i [|\tau(\mathcal{K}_i)^*| \beta_i] \right)^{\frac{1}{r}} \left( \sum_i |\mathcal{K}_i^*| \right)^{\frac{1}{s}} \leq \kappa |J_0|$$

(with the obvious modification for  $q = q_*$ ) and

$$\begin{aligned}
\|A_1 a\|_{H_Y^1} &\leq \frac{p}{p-1} \kappa^{\frac{1}{q_*}} \|A_p\| |J_0|^{\frac{1}{q_*}} \left( \sum_i \|a_i\|_{L_X^{p_*}}^{p_*} \right)^{\frac{1}{p_*}} \\
&\leq \frac{p}{p-1} \kappa^{1+\frac{1}{q_*}} \|A_p\| |J_0|^{\frac{1}{q_*}} \|a\|_{L_X^{p_*}} \\
&\leq \frac{p}{p-1} \kappa^{1+\frac{1}{q_*}} \|A_p\| |J_0| \|a\|_{L_X^\infty} \\
&\leq \frac{p}{p-1} \kappa^{1+\frac{1}{q_*}} \|A_p\|.
\end{aligned}$$

It is not difficult to check that any atom  $a \in H_X^{1,at}(\mathcal{D}_0^N)$  can be written as finite convex combination of atoms considered in this proof so far. Using this and (2) we end up with

$$\|A_1 a\|_{H_Y^1} \leq \frac{p}{p-1} \kappa^{1+\frac{1}{q_*}} \|A_p\| \|a\|_{H_X^{1,at}} \leq \frac{18p}{p-1} \kappa^{1+\frac{1}{q_*}} \|A_p\| \|a\|_{H_X^1}$$

for all  $a \in H_X^1(\mathcal{D}_0^N)$ . ■

Now we interpolate between  $H^1$  and  $L^p$ :

**Lemma 5.5.** *Let  $1 < q < p < \infty$  and  $\frac{1}{q} = \frac{1-\theta}{1} + \frac{\theta}{p}$ . If  $Y$  is a UMD-space, then one has*

$$\begin{aligned} \|A_q : L_X^q(\mathcal{D}_0^N) \rightarrow L_Y^q(\mathcal{D}_0^L)\| \\ \leq c \|A_1 : H_X^1(\mathcal{D}_0^N) \rightarrow H_Y^1(\mathcal{D}_0^L)\|^{1-\theta} \|A_p : L_X^p(\mathcal{D}_0^N) \rightarrow L_Y^p(\mathcal{D}_0^L)\|^\theta \end{aligned}$$

where  $c > 0$  depends at most on  $Y$ ,  $p$ , and  $q$ . In the case  $\gamma_I \equiv 1$  the UMD-property of  $Y$  is not needed and  $c > 0$  does not depend on  $Y$ .

**PROOF.** Because we work with probability spaces consisting of a finite number of atoms only, we can replace (for simplicity)  $X$  and  $Y$  by finite dimensional subspaces  $E \subseteq X$  and  $F \subseteq Y$  such that  $S(E) \subseteq F$ , where we will see that the constant  $c$  can be chosen uniformly for all subspaces  $E$  and  $F$ . The family  $(A_q)_{q \in [1,p]}$  is embedded into an analytic family of operators. Let  $V$  denote the vertical strip  $V = \{x + it : x \in (0, 1), t \in \mathbb{R}\}$  and let

$$J_z(a) := \sum_{I \in \mathcal{D}_0^N} S a_I \gamma_I^{1-z(1-\frac{1}{p})} h_{\tau(I)}.$$

As  $\frac{1}{q} = \frac{1-\theta}{1} + \frac{\theta}{p}$  we have

$$J_\theta = A_q.$$

Since

$$\Re \left( 1 - it \left( 1 - \frac{1}{p} \right) \right) = 1 \quad \text{and} \quad \Re \left( 1 - (1 + it) \left( 1 - \frac{1}{p} \right) \right) = \frac{1}{p},$$

we have

$$\|J_{1+it}(f)\|_{L_F^p(\mathcal{D}_0^L)} \leq 2\text{UMD}_p(Y) \|A_p(f)\|_{L_F^p(\mathcal{D}_0^L)} \quad (6)$$

and

$$\|J_{it}(f)\|_{H_F^1(\mathcal{D}_0^L)} \leq c \|A_1(f)\|_{H_F^1(\mathcal{D}_0^L)} \quad (7)$$

for some  $c > 0$  depending on  $Y$  only. The latter estimate ( $Y$  is a UMD-space) is folklore and can be derived in various ways. For example, one can follow [9, Remarque 2]. Following the proof that the complex interpolation method with parameter  $\theta$  yields an exact interpolation functor of exponent  $\theta$ , for example presented in [1, Theorem 4.1.2], we get that

$$\begin{aligned} & \|J_\theta(f)\|_{(H_F^1(\mathcal{D}_0^L), L_F^p(\mathcal{D}_0^L))_\theta} \\ & \leq \sup_{t \in \mathbb{R}} \|J_{it} : H_E^1(\mathcal{D}_0^N) \rightarrow H_F^1(\mathcal{D}_0^L)\|^{1-\theta} \sup_{t \in \mathbb{R}} \|J_{1+it} : L_E^p(\mathcal{D}_0^N) \rightarrow L_F^p(\mathcal{D}_0^L)\|^\theta \\ & \leq c^{1-\theta} (2\text{UMD}_p(Y))^\theta \|A_1 : H_E^1(\mathcal{D}_0^N) \rightarrow H_F^1(\mathcal{D}_0^L)\|^{1-\theta} \\ & \quad \|A_p : L_E^p(\mathcal{D}_0^N) \rightarrow L_F^p(\mathcal{D}_0^L)\|^\theta \|f\|_{(H_E^1(\mathcal{D}_0^N), L_E^p(\mathcal{D}_0^N))_\theta} \end{aligned}$$

where  $(Z_0, Z_1)_\theta$  denotes the interpolation space obtained by the complex method as in [1, p. 88]. Using

$$(H_E^1(\mathcal{D}_0^N), L_E^p(\mathcal{D}_0^N))_\theta = L_E^q(\mathcal{D}_0^N) \quad \text{and} \quad (H_F^1(\mathcal{D}_0^L), L_F^p(\mathcal{D}_0^L))_\theta = L_F^q(\mathcal{D}_0^L) \quad (8)$$

with multiplicative constants not depending on  $(N, L, X, Y)$  we arrive at our assertion. In the case  $\gamma_I = 1$  we have  $J_{it} = A_1$  and  $J_{1+it} = A_p$  so that the UMD-property in (6) and (7) is not needed. The equivalences (8) are folklore, see [3, p. 334]. One can deduce them via the real interpolation method by exploiting  $(H_Z^1(\mathcal{D}_0^M), L_Z^r(\mathcal{D}_0^M))_{\eta,s} = L_Z^s(\mathcal{D}_0^M)$  for  $\eta \in (0, 1)$ ,  $r, s \in (1, \infty)$  with  $(1/s) = 1 - \eta + (\eta/r)$ ,  $Z \in \{E, F\}$ , and  $M \geq 0$ , where the multiplicative constants in the norm estimates depend on  $(\eta, r, s)$  only (see [16] and the references therein), and the connection between the real and complex interpolation method presented in the second statement of [1, Theorem 4.7.2], where we use that the proof for the first inclusion works as well with  $\theta_0 = 0$ ,  $p_0 = 1$ , and  $(\bar{A})_{\theta_0, p_0}$  replaced by  $A_0$ . ■

**Corollary 5.6.** *Let  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  be a bijection such that there is an  $A \geq 1$  with*

$$\frac{1}{A} \llbracket \mathcal{E} \rrbracket \leq \llbracket \tau(\mathcal{E}) \rrbracket \leq A \llbracket \mathcal{E} \rrbracket \quad (9)$$

*for all non-empty  $\mathcal{E} \subseteq \mathcal{D}$ . Furthermore, let  $X$  be a UMD-space,  $\gamma_I := |I|/|\tau(I)|$ , and  $1 < q < p < \infty$ . Then the boundedness of*

$$\text{Id}_X \otimes T_{p,\tau} : L_{X,0}^p \rightarrow L_X^p$$

*implies the boundedness of*

$$\text{Id}_X \otimes T_{q,\tau} : L_{X,0}^q \rightarrow L_X^q.$$

*In case of  $|\tau(I)| = |I|$  the UMD-property is not needed.*

PROOF. (a) For all  $N \geq 0$  we choose  $L_N \geq 0$  such that

$$\tau(\mathcal{D}_0^N) \subseteq \mathcal{D}_0^{L_N}.$$

Then we can consider the restrictions  $\tau_N : \mathcal{D}_0^N \rightarrow \mathcal{D}_0^{L_N}$  for  $N \geq 0$ . According to Example 5.3 the property  $C(X, p, \kappa)$  for some  $\kappa > 0$  is satisfied uniformly in  $N$ . Applying Lemma 5.5 and Theorem 5.4 gives that

$$\begin{aligned} & \|T_{q,\tau_N} : L_X^q(\mathcal{D}_0^N) \rightarrow L_X^q(\mathcal{D}_0^{L_N})\| \\ & \leq c_{(5.5)} \|T_{1,\tau_N} : H_X^1(\mathcal{D}_0^N) \rightarrow H_X^1(\mathcal{D}_0^{L_N})\|^{1-\theta} \\ & \quad \|T_{p,\tau_N} : L_X^p(\mathcal{D}_0^N) \rightarrow L_X^p(\mathcal{D}_0^{L_N})\|^\theta \\ & \leq c_{(5.5)} \left( \frac{18p}{p-1} \kappa^{1+\frac{1}{q_*}} \right)^{1-\theta} \|T_{p,\tau_N} : L_X^p(\mathcal{D}_0^N) \rightarrow L_X^p(\mathcal{D}_0^{L_N})\| \\ & =: c \|T_{p,\tau_N} : L_X^p(\mathcal{D}_0^N) \rightarrow L_X^p(\mathcal{D}_0^{L_N})\| \\ & \leq c \|T_{p,\tau} : L_{X,0}^p \rightarrow L_X^p\|. \end{aligned}$$

(b) Now we consider a strictly increasing sequence of integers  $B_N \geq 1$  such that

$$\tau(\mathcal{D}_0^{B_N}) \supseteq \mathcal{D}_0^N.$$

For  $a = \sum_{I \in \mathcal{D}} a_I h_I$ , where  $(a_I)_{I \in \mathcal{D}} \subseteq X$  is finitely supported, we get

$$\|T_{q,\tau} a\|_{L_X^q} = \sup_N \|E(T_{q,\tau} a | \mathcal{F}_N)\|_{L_X^q}$$

$$\begin{aligned}
&= \sup_N \|E(T_{q,\tau_{B_N}} a_{B_N} | \mathcal{F}_N)\|_{L_X^q} \\
&\leq \sup_N \|T_{q,\tau_{B_N}} a_{B_N}\|_{L_X^q} \\
&\leq \sup_N \|T_{q,\tau_{B_N}} : L_X^q(\mathcal{D}_0^{B_N}) \rightarrow L_X^q(\mathcal{D}_0^{L_{B_N}})\| \|a_{B_N}\|_{L_X^q(\mathcal{D}_0^{B_N})} \\
&\leq c \|T_{p,\tau} : L_{X,0}^p \rightarrow L_X^p\| \|a\|_{L_{X,0}^q}
\end{aligned}$$

where  $\tau_{B_N} : \mathcal{D}_0^{B_N} \rightarrow \mathcal{D}_0^{L_{B_N}}$  is the restriction of  $\tau$  considered in (a) and  $a_{B_N}$  the restriction of  $a$  to  $\mathcal{D}_0^{B_N}$ . ■

Modifying slightly the first step in the proof of Corollary 5.6 we can remove the assumption that  $X$  is a UMD-space in Corollary 4.3:

**Corollary 5.7.** *Let  $X$  be a Banach space and let  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  be a permutation such that  $|\tau(I)| = |I|$ . Then, for  $1 < q < p < 2$ , the boundedness of*

$$\text{Id}_X \otimes T_\tau : L_{X,0}^p \rightarrow L_X^p$$

*implies the boundedness of  $\text{Id}_X \otimes T_\tau : L_{X,0}^q \rightarrow L_X^q$ .*

PROOF. Our assumption implies  $\gamma_I = 1$  and that  $\tau$  satisfies Semenov's condition with some  $\kappa \in [1, \infty)$ . By Example 5.2 the restrictions  $\tau_N : \mathcal{D}_0^N \rightarrow \mathcal{D}_0^N$  satisfy condition  $c(X, p, \kappa)$  for all  $p \in (1, \infty)$ . Now we can follow the proof of Corollary 5.6 with  $L_N = B_N = N$  and  $\gamma_I = 1$  so that the UMD-property in Lemma 5.5 is not needed. ■

We close with an extrapolation theorem for rearrangement operators that are isomorphisms on  $L_{X,0}^p$ . For real valued rearrangements, i.e. when  $X = \mathbb{R}$ , the following theorem is well known. It can be obtained by different methods, the most direct route [10] going via Pisier's re-norming in  $L^p$ .

**Theorem 5.8.** *Let  $\tau : \mathcal{D} \rightarrow \mathcal{D}$  be a bijection and  $\gamma_I := |I|/|\tau(I)|$ . Assume that  $X$  is a UMD-space. If there exists a  $p \in (1, \infty)$  with  $p \neq 2$  such that*

$$\|\text{Id}_X \otimes T_{p,\tau} : L_{X,0}^p \rightarrow L_X^p\| \cdot \|\text{Id}_X \otimes T_{p,\tau^{-1}} : L_{X,0}^p \rightarrow L_X^p\| < \infty, \quad (10)$$

*then for each  $q \in (1, \infty)$  one has that*

$$\|\text{Id}_X \otimes T_{q,\tau} : L_{X,0}^q \rightarrow L_X^q\| \cdot \|\text{Id}_X \otimes T_{q,\tau^{-1}} : L_{X,0}^q \rightarrow L_X^q\| < \infty. \quad (11)$$

PROOF. (a) First we observe that our assumption implies that (10) holds for  $X = \mathbb{C}$  and  $X = \mathbb{R}$ . If  $p \in (2, \infty)$ , then [11, Theorems 2 and 3] imply condition (9). In case of  $p \in (1, 2)$  duality implies (10) for  $X = \mathbb{R}$  and  $p$  replaced by the conjugate index  $p' \in (2, \infty)$ . Hence we have (9) as well.

(b) From Corollary 5.6 and (a) we immediately get (11) for  $q \in (1, p)$ .

(c) Let  $q \in (p, \infty)$ . It is easy to see that for a bijection  $\sigma : \mathcal{D} \rightarrow \mathcal{D}$  and  $r \in (1, \infty)$  the boundedness of

$$\| \text{Id}_X \otimes T_{r,\sigma} : L_{X,0}^r \rightarrow L_X^r \| \quad \text{and} \quad \| \text{Id}_{X'} \otimes T_{r',\sigma^{-1}} : L_{X',0}^{r'} \rightarrow L_{X'}^{r'} \|$$

are equivalent to each other where  $1 = (1/r) + (1/r')$  (note, that  $X$  is in particular reflexive because of the UMD-property). Using this observation our assumption (10) holds for  $p'$  and  $X'$  and the conclusion for  $q' \in (1, p')$  and  $X'$ . By duality we come back to  $q$  and  $X$ . ■

## References

- [1] Bergh, J. and Löfström, J., Interpolation spaces. An Introduction. Springer 1976.
- [2] Bernard, A. and Maisonneuve, B., Decomposition atomique de martingales de la class  $H_1$ . *Sem. Prob. XI*, Lecture Notes Math. 581:303–323, 1977.
- [3] Blasco, O. and Xu, Q., Interpolation between vector valued Hardy spaces. *J. Funct. Anal.*, 102(2):331–359, 1991.
- [4] Burkholder, D.L., Explorations in martingale theory and its applications. Ecole d'Été de Probabilités de Saint-Flour, XIX–1989, *Lect. Notes Math.* 1464:1–66, 1992, Springer.
- [5] Burkholder, D.L., Martingales and singular integrals in Banach spaces. In: *Handbook of the geometry of Banach spaces*, Vol. I, 233–269, North-Holland, Amsterdam, 2001.
- [6] Burkholder, D.L. and Gundy R.F., Extrapolation and interpolation of quasilinear operators on martingales. *Acta Math.* 124: 249–304, 1970.



- [7] Coifman, R.R., A real variable characterization of  $H^p$ . *Studia Math.* 51:269–274, 1974.
- [8] Geiss, S. , Müller, P. F. X. and Pillwein, V., A remark on extrapolation of rearrangement operators on dyadic  $H^s$ ,  $0 < s \leq 1$ . *Studia Math.*, 171:197–205, 2005.
- [9] B. Maurey, Système de Haar. *Seminaire Maurey–Schwartz, Ecole Polytechnique, Paris*, 1974–1975.
- [10] Müller, P. F. X., Isomorphisms between  $H^1$  spaces. Birkhäuser Verlag, Basel, 2005.
- [11] Müller, P. F. X., Rearrangements of the Haar system that preserve BMO. *Proc. London Math. Soc. (3)*, 75(3):600–618, 1997.
- [12] Müller, P. F. X. and Schechtman, G., Several results concerning unconditionality in vector valued  $L^p$  and  $H^1$  spaces. *Illinois J. Math.*, 35:220–233, 1991.
- [13] Semenov, E. M., Equivalence in  $L^p$  of permutations of the Haar system. *Dokl. Akad. Nauk SSSR*, 242(6):1258–1260, 1978.
- [14] Semenov, E. M. and Stöckert, B., The rearrangements of the Haar system in the spaces  $L_p$ . *Anal. Mathematica*, 7:277–295, 1981.
- [15] Weisz, F., Martingale Hardy spaces for  $0 < p < 1$ . *Prob. Theory Rel. Fields*, 84:361–376, 1990.
- [16] Weisz, F., Martingale operators and Hardy spaces generated by them. *Studia Math.*, 114:39–70, 1995.

### Addresses

Department of Mathematics and Statistics  
P.O. Box 35 (MaD)  
FIN-40014 University of Jyväskylä  
Finland

Department of Analysis  
J. Kepler University  
A-4040 Linz  
Austria