A correction to: Sobolev inequalities on sets with irregular boundaries

by Tero Kilpeläinen and Jan Malý

February 12, 2008

The proof of Theorem 2.3 in the original paper that appeared in Z. Anal. Anwendungen vol 19 (2000), no. 2, 369–380, was slightly erroneous. Here we provide a new proof:

Theorem 2.3 *Suppose that Ω is an s-John domain and $b \geq 1-n$. Then there is a constant $C = C(n,p,q,\Omega) > 0$ such that*

$$\Big(\int_\Omega |u-\bar{u}_a|^q \rho^a \, dx\Big)^{1/q} \leq C\Big(\int_\Omega |\nabla u|^p \rho^b dx\Big)^{1/p}$$

for each $u \in C^1(\Omega)$; here the Sobolev exponent is

$$q = \frac{p(n+a)}{s(n+b-1)-p+1}.$$

Proof. We will verify the assumptions of Theorem 2.1. First we notice that the inequalities $s \geq 1$ and $b \geq 1-n$ imply

$$p\Big(\frac{a+n}{q} + 1 - \frac{n}{p}\Big) = s(n+b-1)+1-n \geq b,$$

so that (2.4) is true. For fixed $y \in \Omega \setminus B_0$, the s-John core γ on $[0,\ell]$ gives us the desired worm: Let m be the integer such that

$$3\ell < 2^m \rho(y) \leq 6\ell.$$

Since

$$\rho(y) \le \rho(x_0) + |y - x_0| \le 3|y - x_0| \le 3\ell,$$

we have $m \ge 1$. Set

$$\xi_k = 2^{k-m}\ell.$$

Then $(\gamma, \{\xi_k\})$ is a worm with parameters m, $\{\ell_k\}$, $\{R_k\}$, $\{r_k\}$. Observe that we may assume that the part of γ lying inside the ball $B(y, 2^{1-m}\ell) \subset \Omega$ is a line segment. Thus we have that

$$\rho(y) \le 6\, 2^{-m}\ell \le 3R_1 \le 3R_k$$

so that (2.5) holds.

Moreover,

$$\begin{aligned} &\ell_k \le \xi_k, \\ &\xi_k \ge R_k, \\ &r_k \ge c_0 \xi_{k-1}^s = c_0(\xi_k/2)^s. \end{aligned}$$

Since

$$(n+a)/q = (s(n+b-1)+1-p)/p$$

we have by choosing $\tau_k = \min(2^{k-m}, \xi_k)^{(n+a)/q} \in (0,1]$ that

$$\tau_k \le \xi_k^{(n+a)/q} \le C\tau_k\,,$$

where the constant C does not depend on y since for the s-John core γ it holds that

$$c_0\ell^s \le \operatorname{diam}(\Omega).$$

Thus we have

$$\boldsymbol{\mu}(B(y, 3R_k))^{1/q} \le C\xi_k^{(n+a)/q} \le C\tau_k$$

and furthermore,

$$\begin{aligned} r_k^{-(n+b-1)/p}\ell_k^{(p-1)/p} &\le C\big(\xi_k\big)^{-s(n+b-1)/p}\xi_k^{(p-1)/p} \\ &= C\xi_k^{-(n+a)/q} \\ &\le C\tau_k^{-1}\,. \end{aligned}$$

Hence the claim follows from Theorem 2.1.